

# PHOSPHOLIPASE A<sub>2</sub> (PLA<sub>2</sub>) REGULATES NEUROEXOCYTOSIS TO COUNTERACT BOTULINUM TOXIN A (BoNT/A) POISONING

Xieping Zhang, Hiroshi Ishida and Prabhati Ray

Section on Molecular Biology

Department of Biology, Division of ET

Walter Reed Army Institute of Research, Silver Spring, MD 20910

## ABSTRACT

In this study, BoNT/A is one of the most serious biological threats faced by the U. S. military and allied forces. Previously we reported that in nerve growth factor (NGF)-differentiated PC12 cells, arachidonic acid (AA) release is associated with acetylcholine (ACh) release and BoNT/A inhibits both. We report the effect of PLA<sub>2</sub> over expression on inhibition of ACh exocytosis due to BoNT/A light chain (LC) in PC12 cells. Over expression of PLA<sub>2</sub> alone augmented the stimulated release of ACh and AA. PLA<sub>2</sub> over expression also effectively prevented the inhibition of stimulated ACh and AA release due to BoNT/A LC.

## INTRODUCTION

Botulinum toxin type A (BoNT/A) acts by blocking Ca<sup>2+</sup>-dependent ACh release (neuroexocytosis) at peripheral neuromuscular junctions, sometimes causing fatal neuromuscular paralysis. The clonal pheochromocytoma PC12 cell line was used to study the biochemical mechanisms of action of BoNT/A. To explain the mechanism of action of BoNT/A, we previously reported that when NGF differentiated PC12 cells are stimulated with high K<sup>+</sup> (80 mM), AA and ACh are co-released and BoNT/A inhibits both release processes. (Ray, P. *et al.*, 1993, J. Biol. Chem., 268, 11057-11064). Stimulus-induced ACh release and AA release are inhibited in the presence of Ca<sup>2+</sup>-dependent PLA<sub>2</sub> inhibitors (AACOCF<sub>3</sub>, DEDA). On the contrary, PLA<sub>2</sub> activators such as mastoparan can cause ACh release in cells (Ray, P. *et al.*, 1997, NeuroReport, 8, 2271-2274).

PLA<sub>2</sub> acts on membrane phospholipids to generate AA and lysophosphatidic acid (LPA). AA acts as a fusogen to induce exocytosis. LPA activates Rho GTPases, which have been implicated in mechanisms correlating actin cytoskeletal organization and exocytosis. In this study, we show that PLA<sub>2</sub> over expression prevented the inhibition of stimulated ACh and AA release due to BoNT/A LC.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>01 JUL 2003</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Phospholipase A2 (Pla2) Regulates Neuroexocytosis To Counteract Botulinum Toxin A (Bont/A) Poisoning</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Section on Molecular Biology Department of Biology, Division of ET Walter Reed Army Institute of Research, Silver Spring, MD 20910</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM001523., The original document contains color images.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>5</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## METHODS

### CELL CULTURE

PC12 cells (CLONTECH) were grown in 75-cm<sup>2</sup> tissue culture flasks in DMEM (GIBCO) containing 5 % (v/v) FBS 10 % (v/v) donor horse serum (CLONTECH), streptomycin 100 µg/ml and 100 units/ml penicillin G sodium (Sigma). The cells ( $1.5 \times 10^6$ ) were incubated at 37°C in a humidified atmosphere of 90% air 10% CO<sub>2</sub>. After 3 days the cells were differentiated with nerve growth factor (NGF, 50 ng/ml growth medium) for 4 days.

### sPLA<sub>2</sub> OVER EXPRESSION IN PC12 CELLS

The levels of sPLA<sub>2</sub> in differentiated PC12 cells after transfection with pTracer-PLA<sub>2</sub> was analysed by immunoprecipitation and Western blotting analysis using rabbit antiserum against sPLA<sub>2</sub> (Cayman Chemical ).

### BoNT/A LC OVER EXPRESSION IN PC12 CELLS

Expression of BoNT/A LC after transfection was analysed by Western blotting with use of GFP monoclonal antibody (CLONTECH).

### SNAP-25 DETECTION

SNAP-25 was visualized by SNAP-25 antibody and TRITC conjugated IgG.

### ARACHIDONIC ACID (AA) RELEASE ASSAY

Arachidonic acid release assay was performed by modification of the method of Ray.P et al. as described (J. Biol. Chem., 268, 11057-11064).

### ACETYLCHOLINE (ACH) RELEASE ASSAY

Acetylcholine release assay was performed by modification of method of Ray.P et al. as described (J. Biol. Chem., 268, 11057-11064).

## RESULTS




Figure 1. Transfection efficiency of PC 12 cells by pEGFP-BoNT/A LC

The transfection efficiency (>70%) was calculated twenty-four hours and forty-eight hours after transfection with pEGFP-BoNT/A LC by fluorescence microscope. Images were taken twenty-four hours and forty-eight hours after transfection




Figure 2. BoNT/A LC over expression in PC12 cells

Twenty-four hours after transfection of differentiated PC 12 cells with pEGFP-BoNT/A LC, the expression level of LC was confirmed by Western blotting using of GFP antibody. Control: No GFP was expressed; Transfection with pEGFP vector alone: GFP (28 kDa) was expressed; Transfection with pEGFP-BoNT/A LC: GFP-fused LC (78 kDa) was expressed.




Figure 3. Effect of transfection of BoNT/A LC on SNAP-25 degradation

Twenty-four hours after transfection of differentiated PC12 cells with pEGFP-BoNT/A LC, SNAP-25 was visualized by SNAP-25 antibody and TRITC conjugated IgG. There was a marked degradation of SNAP-25 (red) in LC expressing cells (green), but no degradation was seen in untransfected cells.



Figure 4. sPLA<sub>2</sub> over expression

Immunoprecipitation followed by Western blotting analysis with sPLA<sub>2</sub> antibody, showed that transfection with sPLA<sub>2</sub> over expressed sPLA<sub>2</sub> (14 kDa). However, control cell (no transfection) or vector alone transfection did not express detectable amount of the protein.




Figure 5. Effect of cotransfection of BoNT/A LC and PLA<sub>2</sub> on AA and ACh release

NGF-differentiated PC12 cells were transiently transfected with either pEGFP (pEV)-BoNT/A LC or cotransfected with pTracer (pTV)-sPLA<sub>2</sub>. Over expression of PLA<sub>2</sub> in these BoNT/A LC transfected cells effectively prevented the inhibition of stimulated (by high K<sup>+</sup>) ACh and AA release due to BoNT/A LC.

## CONCLUSION

- PLA<sub>2</sub> mediated mechanisms may regulate presynaptic ACh exocytosis.
- BoNT/A inhibits neuroexocytosis by targeting these mechanisms.
- Manipulating PLA<sub>2</sub> mechanisms in neuroexocytosis may serve as prospective strategies to counteract BoNT/A poisoning.

## REFERENCES

1. Ray, P., C. B. Millard, J. P. Petrali, J. D. Berman, and R. Ray. 1997. Acetylcholine exocytosis in PC12 cells deficient in SNAP-25. *NeuroReport* **8**: 2271-2274.
2. Ray, P., J. D. Berman, W. Middleton, and J. Brendle. 1993. Botulinum toxin inhibits arachidonic acid release associated with acetylcholine release from PC12 cells. *J. Biol. Chem.* **268**: 11057-11064.
3. Green L. A. and Tischler, A. S. 1982. PC12 pheochromocytoma cells in neurobiological research. *Adv. Cell. Neurobiol.*, **3**, 373-414.